

Delivery of a nematocyst toxin

SIR—Cnidaria (hydras, jellyfish, sea anemones and corals) possess a wide array of neurotoxic, cytolytic and enzymatic¹ substances. Despite the fact that these toxic substances were isolated mainly from homogenates of entire animals or tentacles, their storage and delivery sites are attributed to nematocysts, the stinging subcellular organelles. Nematocysts consist of a capsule containing a highly folded eversible tubule². The discharge of the nematocysts is driven by

the capsule's high internal hydrostatic pressure of 15 megapascals (≈ 150 atmospheres), which causes the eversion of the tubule with accelerations up to 40,000g and therefore comprises one of the fastest events in biology^{3,4}. However, the ultrastructural and mechanical data do not explain toxin allocation and the mode of their delivery. Further, even the notion that nematocysts serve as the venom source was not unequivocally established.

The present study aimed to establish the chemical nature of jellyfish toxins, their subcellular localization and the route of their delivery. We show that cnidarian venom is stored in nematocysts on the outer surface of the tubule and is

delivered upon its discharge (eversion–extension) through the spirally arranged array of the extended hollow tubular barbs, thus resembling a multiheaded poisonous arrow.

We have purified a phospholipase A₂ toxin (M_r 16,000; ref. 5) from the tentacles of the Mediterranean medusa *Rhopilema nomadica*. Purity was assessed by electrophoresis (*a* in the figure), amino-acid analysis and by amino-terminal amino-acid sequence determination (*g* in the figure). Polyclonal antitoxin antibodies were raised in rabbit⁶, affinity-purified by slot immunoblotting (*b*, *d*) and used as immunocytochemical tools to study toxin allocation by light (*e*, *f*) and electron microscopy⁷. Immunoblot revealed the presence of the toxin in the fishing tentacles, but not in the bell tissues (*b–d*). The bell, in contrast to the tentacles, is devoid of nematocysts. The toxin may therefore be attributed to the venomous hunting apparatus of the jellyfish, and is not likely to be a widely distributed defence chemical.

Immunocytochemical staining of cryostat-sliced nematocysts revealed a highly specific allocation of the toxin limited to their tubular segments (*e*, *f*). In the resting nematocyst (*e*) the capsular lumen and wall were devoid of immunofluorescent staining. The allocation of the

toxin on the discharged tubule revealed a helical shape (*f*) which parallels the helical arrangement of the barbs. The latter observation has raised the possibility that the hollow barbs⁸ may function as a device for toxin delivery. This essential aspect was further studied by high-resolution immunogold technique and electron microscopy⁷. It has been shown that the toxin is externally allocated on the surface of the inverted–undischarged tubule. In the everted–discharged tubule the toxin occurs on its internal surface and accumulates in the vicinity of the bases of the hollow barbs and their lumina (data not shown).

The above data provide a visualization of the route of toxin translocation and release schematically presented in *h*. In the resting nematocyst, toxin is located in the crypts (invaginations) of the outer surface of the twisted, folded and inverted tubule, the lumen of which is occupied by the internalized barbs. During discharge and tubule eversion toxin is translocated from the outer to the inner surface of the everted and extended tubule while aggregating near the bases of and within the hollow barbs during release.

We assume that the hydrostatic pressure raised in the dischargeable capsule⁹, causing the tubule eversion, also supplies the appropriate propulsion of toxin delivery through the barb system. This notion is supported by a study showing that the fibre-like structure of the capsule's inner wall, which provides the tensile strength necessary to withstand the capsule's high osmotic pressure, continues along the tubule's wall³. Thus, the nematocyst delivery system may provide an example of subcellular protein translocation based on simple mechanical forces.

A. Lotan^{††}

L. Fishman[†]

Y. Loya[†]

E. Zlotkin^{*}

^{*}Department of Cell and Animal Biology,

Life Science Institute,

Hebrew University,

Jerusalem 91904, Israel


[†]Department of Zoology,

Faculty of Life Sciences,

Tel Aviv University,

Tel Aviv 69978, Israel

- Walker, M. J. A. in *Marine Toxins and Venoms* (ed. Tu, A. T.) 279–325 (Dekker, New York, 1988).
- Mariscal, R. N. in *Coelenterate Biology* (eds Muscatine, L. & Lenhoff, H. M.) 129–178 (Academic, New York, 1974).
- Holstein, T. W. *et al.* *Science* **265**, 402 (1994).
- Holstein, T. W. & Tardent, P. *Science* **223**, 830 (1984).
- Gomez, F. *et al.* *Eur. J. Biochem.* **186**, 233–33 (1989).
- Vaitukaitis, J. L. *Meth. Enzym.* **73**, 46–52 (1981).
- Castel, M., Belenky, M., Cohen, S., Otterson, O. & Mathisen, J. S. *Eur. J. Neurosci.* **5**, 368 (1993).
- Hessinger, D. A. & Ford, M. T. in *The Biology of Nematocysts* (eds Hessinger, D. A. & Lenhoff, H. M.) 75–94 (Academic, Orlando, 1988).
- Tardent, P. in *The Biology of Nematocysts* (eds Hessinger, D. A. & Lenhoff, H. M.) 309–332 (Academic, Orlando, 1988).



Chemistry, location and delivery of a nematocyst toxin derived from *R. nomadica*. **a**, SDS–PAGE electrophoresis of 2 μ g toxin. **b–d**, Immunoblots. Separation of 10 μ g proteins extracted from the fishing tentacles (*b*), bell tissues (*c*) and 0.1 μ g of purified toxin (*d*). Tentacles of the freshly collected jellyfish were cut at sea, stored over dried ice, then deep-frozen (-80°C). Tentacles were thawed at 25°C followed by centrifugation ($15,000g$, 30 min). The supernatant-containing material from discharged nematocysts was desalted by dialysis and separated by DEAE–cellulose anion-exchange column followed by Mono-S–HPLC cation-exchange column, resulting in a substance toxic to fishes by injection ($\text{LD}_{50} = 0.6 \mu\text{g}$ per 100 mg body weight). Purity of the toxic component was assessed by electrophoresis, amino-acid analysis and N-terminal amino-acid sequence determination. Proteins extracted from jellyfish tentacles were separated on a BioRad minigel apparatus, transferred to nitrocellulose membranes overlaid with the toxin-specific antibodies and detected by chemiluminescence. The western blotting kit (rabbit) was purchased from Boehringer-Mannheim. **e**, **f**, Immunofluorescent light microscopy of a resting (*e*) and discharged (*f*) nematocyst. Note that the stain revealing the toxin occurs in the tubule (*e*) and is limited to its helically arranged dots (*f*) when discharged. Bars: *e*, 2 μm ; *f*, 1 μm . For light microscopy, tentacles were sliced by a Jung CM 3000 (Leica) cryostat to sections of 8 μm , attached to glass slides with polylysine, incubated with the affinity-purified antitoxin antibody, then with goat anti-rabbit fluorescein isothiocyanate-labelled antibody (Jackson Inc.). **g**, N-terminal amino-acid sequence of the new toxin. This segment has about 60% amino-acid homology with toxic phospholipases A₂ derived from the venom of the *Heloderma* lizard and a honey bee⁵. **h**, Schematic presentation of toxin compartmentation and delivery in the nematocyst system. (1), Resting tubule, whose lumen is filled with barbs. The toxin (diamonds) is located in the folds and invaginations of the tubule's membrane (black line). (2), During discharge, toxin is translocated into the tubule while the barbs emerge and extend. (3), Toxin delivered from the everted–extended tubule through the hollow barbs.